ΑD)	

Award Number: W81XWH-10-1-0662

TITLE: SIRT3 Is a Mitochondrial Tumor Suppressor and Genetic Loss Results in a Murine Model for ER/PR-Positive Mammary Tumors Connecting Metabolism and Carcinogenesis

PRINCIPAL INVESTIGATOR: Sarki Abdulkadir

CONTRACTING ORGANIZATION: Vanderbilt University Medical Center Nashville, TN 37203-6876

REPORT DATE: September 2012

TYPE OF REPORT: Revised Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1 September 2012	Revised Annual	1 September 2011 - 31 August 2012
4. TITLE AND SUBTITLE	Tromod / timedi	5a. CONTRACT NUMBER
SIRT3 Is a Mitochondrial Tumor Su	5b. GRANT NUMBER	
Model for ER/PR-Positive Mammary	W81XWH-10-1-0662	
Carcinogenesis	5c. PROGRAM ELEMENT NUMBER	
•		
6. AUTHOR(S)		5d. PROJECT NUMBER
Sarki Abdulkadir		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
E-Mail: sarki.abdulkadir@vanderbilt.e		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT NUMBER
Vanderbilt University Medical Cente	or .	NUMBER
Nashville, TN 37203-6876		
Nasilville, 114 37203-0070		
9. SPONSORING / MONITORING AGENCY	NAME(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)
U.S. Army Medical Research and M		is a creatiment of terrain (e)
Fort Detrick, Maryland 21702-5012		
		11. SPONSOR/MONITOR'S REPORT
		NUMBER(S)
		`,
40 DIOTRIBUTION / AVAIL ABILITY OTATI	-14-11	

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for Public Release; Distribution Unlimited

13. SUPPLEMENTARY NOTES

14. ABSTRACT

Vj g'i qcnı'qh'y ku'u{pgti kuke'r tqlgev'ctg'\q'guvcdrkuj "yj g'o kqej qpf tkc''nqecrk gf "uktwkp''r tqvgkp''UKTV5''cu'c''wo qt"
uwr r tguuqt "kp''dtgcuv'ecpegt "cpf "\q'f ghkpg''ku''tqrg''cu'c''o qrgewct "rkprn'dgwy ggp"ci kpi "cpf "dtgcuv'ecpegt0'Uktv5''npqenqw''
o keg'f gxgmr 'GT IRT/r qukkxg''dtgcuv''wo qtu''rcvgt "kp''rktg0'Vj gug''GT IRT/r qukkxg''wo qtu''ctg'j kuvqmi kecm{ "uko krct "\q'dtgcuv''
wo qtu''eqo o qp''kp''qrf gt''y qo gp0'Kp''j wo cpu. "nquu''qh''UKTV5''ku''uggp''kp''c''uki pkhkecpv'htcevkqp''qh''dtgcuv'ecpegtu''cpf "o c{"
ugtxg''cu''c''o qrgewct''dkqo ctngt0'O qrgewct''vcti gwu''qh'UKTV5''f gcegv{rcvkqp''j cxg''dggp''kf gpvkhkgf. "kpenwf kpi "O pUQF "cpf"
QUER0Cpvkdqf kgu''yj cv'tgeqi pk g''ur gekhke''cegv{rcvgf "n\ukpg''tgukf wgu''vcti gvgf "d{"UKTV5''kp''yj gug''o qrgewrgu''j cxg''dggp''
kf gpvkhkgf "cpf "xcrkf cvgf "cpf "ctg''dgkpi "f gxgmr gf "cu''r qvgpvkcn'pqxgn''dkqo ctngtu''kp''dtgcuv''ecpegt0'''Vj gug''uwrf kgu''j cxg''
gpj cpegf "qwt''wpf gtuvcpf kpi "qh''yj g'o qrgewct''hpmu''dgwy ggp''ci kpi "cpf "dtgcuv''ecpegt"cpf "r tqxkf g''pqxgn'r qvgpvkcn''
dkqo ctngtu''qh''dtgcuv''ecpegt "kp''j wo cpu0

15. SUBJECT TERMS

Breast cancer, SIRT3, acetylation, aging, biomarkers						
BER 19a. NAME OF RESPONSIBLE PERSON						
S USAMRMC						
19b. TELEPHONE NUMBER (include area						
code)						

Table of Contents

	<u>Page</u>
Introduction	3
BODY	3-5
Key Research Accomplishments	5
Reportable Outcomes	5
Conclusion	5
References	5

INTRODUCTION

The mammalian *Sirtuin* genes are homologs of the yeast Saccharomyces cerevisiae *Sir2* gene that is implicated in the regulation of longevity (Haigis et al., 2012). There are seven Sirtuin proteins, with SIRT1, SIRT6, and SIRT7 localized in the nucleus, SIRT2 localized in the cytoplasm and SIRT3, SIRT4, and SIRT5 localized in the mitochondria (Finkel et al., 2009). The mitochondrial deacetylase SIRT3 is thought to act on numerous substrates to regulate seevral processes including fat and amino-acid metabolism as well as electron transport (Huang et al., 2010).

SIRT3 has been proposed to function in maintaining mitochondrial integrity and to serve as a bonafide tumor suppressor (Finley et al., 2011; Kim et al., 2010). We have shown that Sirt3-/- MEFs expressing only Myc or Ras are able to grow in soft agar and form tumors in nude mice, unlike wild type MEFS which require the expression of both oncogenes (Kim et al., 2010). Furthermore, Sirt3 knockout mice develop estrogen receptor and progesterone receptor (ER/PR) positive breast cancers (Kim et al., 2010).

One third of female Sirt3 knockout mice developed mammary gland tumors by 24 months. These tumors were well-differentiated, ER/PR+ tumors similar to the tumors commonly seen in breast malignancies in older women. Our analysis of 992 human breast cancer samples from human tumor mRNA expression databases showed a significant reduction in SIRT3 mRNA in breast cancers compared to benign tissue as well as an association with grade (Kim et al., 2010).

BODY

Statement of Work - Task 1 - Identify Sirt3 mitochondrial deacetylation targets and determine if these targets are regulated by extracellular stimuli known to activate sirtuin function (e.g., resveratrol). These targets will subsequently be knocked down (with siRNA) to determine if there is a mechanistic connection between the increase in superoxide and the stress-induced genomic instability observed in SIRT3^{-/-} cells (months 1-18).

Results: The results for Task 1 have been detailed in the report for W81XWH-10-1-0661, which is the companion grant for W81XWH-10-1-0662 in this synergistic award mechanism. These studies have identified Manganese Superoxide Dismutase (MnSOD) and OSCP as SIRT3 targets. The specific lysine residues in these proteins targeted by Sirt3 have been identified and new antibodies have been developed. These include the OSCP acetylated K139 and MnSOD acetylated K68 antibodies.

Statement of Work - Task 2 - Determine if exposure to resveratrol or overexpression of a MnSOD gene will prevent increases in ROS in MEFs and/or decrease the development of mammary tumors in *Sirt3* knockout mice and transformation in SIRT3^{-/-} MEFs (months 7-24).

Results: The mice for this study are being generated by W81XWH-10-1-0661 while W81XWH-10-1-0662 will be involved in the analysis of the mammary tumors in Sirt3 knockout mice. The animals have been bred and are being aged, since mammary tumors develop in older mice after 1 year of age. These studies were slated to start in month 7 and are ongoing.

Statement of Work - Task 3 - Determine if loss of SIRT3 ductal protein in ER/PR-positive and - negative breast samples from the Vanderbilt Breast Spore correlates with clinically significant outcomes including response to therapy, local tumor control, disease free survival, and overall survival (months 1-24).

We have used a human breast tissue array to determine conditions for SIRT3 staining (Figure 1). We have validated the new antibodies generated against acetylated SIRT3 substrates, including MnSOD-K68 and OSCP-K138. We described in the first year annual report the western blot analyses validation of these antibodies. We next worked to define the conditions for immunohistochemistry using these antibodies. Because human

breast cancer samples available to us are available as formalin-fixed, paraffin-embedded (FFPE) tissue, we sought to work out antibody conditions for staining FFPE tissue sections. We tested antigen retrieval conditions in citrate buffer, pH 6 and EDTA buffer pH9. Figure 2 shows examples of images from successful staining by the new antibodies in FFPE tissues by immunohistochemistry in Sirt3 knockout mouse tissue. We are now validating these findings in human sections before proceeding to analysis of tissue microarrays.




Figure 1. Examples of SIRT3 staining pattern in a breast cancer tissue array, showing various degrees of loss of SIRT3 staining.




Figure 2. Validation of the OSCP acetylated K139 and MnSOD acetylated K68 antibodies. Sirt3 knockout mouse prostate sections were stained with the indicated antibodies and positive glands showing cytoplasmic staining detected by Nova red staining (arrows). Negative glands on the same sections serve as internal negative control. No primary antibody controls shows no staining (not shown).

This work is on-going and was slightly delayed to accommodate the validation of the new antibodies may be molecular biomarkers. In this task we will determine clinicopathologic variables including age, tumor grade, stage, mitotic count, and ER and PR status and relate these to SIRT3 and its substrates. We will be submitting a no cost extension to allow completion of these studies as outlined in the statement of work.

KEY RESEARCH ACCOMPLISHMENTS:

- 1. Validation of SIRT3 loss in human breast cancer samples.
- 2. Validation of antibodies for SIRT3 lysine substrates in MnSOD and OSCP in immunohistochemistry using formalin-fixed paraffin embedded samples.

REPORTABLE OUTCOMES:

1. Antibodies for SIRT3 lysine substrates in MnSOD and OSCP suitable for immunohistochemistry

CONCLUSION:

Aging has long been recognized as a risk factor for breast cancer incidence, but the molecular basis for this association is not understood. Our studies suggest that SIRT3 may provide a molecular link between breast cancer and aging. This synergistic DOD idea award has enabled the development of a Sirt3 knockout mouse model of aging-related receptor-positive breast cancers. In addition, new potential biomarkers for breast cancer are being developed. These studies provide important new insights into breast tumorigenesis as well as provide possible new therapeutic and prognostic targets for human breast cancer.

REFERENCES

Finkel, T., Deng, C.X., and Mostoslavsky, R. (2009). Recent progress in the biology and physiology of sirtuins. Nature *460*, 587-591.

Finley, L.W., Carracedo, A., Lee, J., Souza, A., Egia, A., Zhang, J., Teruya-Feldstein, J., Moreira, P.I., Cardoso, S.M., Clish, C.B., *et al.* (2011). SIRT3 opposes reprogramming of cancer cell metabolism through HIF1alpha destabilization. Cancer Cell *19*, 416-428.

Haigis, M.C., Deng, C.X., Finley, L.W., Kim, H.S., and Gius, D. (2012). SIRT3 is a mitochondrial tumor suppressor: a scientific tale that connects aberrant cellular ROS, the Warburg effect, and carcinogenesis. Cancer Res 72, 2468-2472.

Huang, J.Y., Hirschey, M.D., Shimazu, T., Ho, L., and Verdin, E. (2010). Mitochondrial sirtuins. Biochim Biophys Acta *1804*, 1645-1651.

Kim, H.S., Patel, K., Muldoon-Jacobs, K., Bisht, K.S., Aykin-Burns, N., Pennington, J.D., van der Meer, R., Nguyen, P., Savage, J., Owens, K.M., *et al.* (2010). SIRT3 is a mitochondria-localized tumor suppressor required for maintenance of mitochondrial integrity and metabolism during stress. Cancer Cell *17*, 41-52.